

**AML 19**  
**Acute Promyelocytic Leukaemia**  
**AIDA Consolidation cycle 2**

<b>Height</b>	cm
<b>Weight</b>	kg
<b>BSA</b>	m <sup>2</sup>

AML 19 patient trial number:

Destination:	BMTU
	HDW

	Limits
Date	
Neutrophils	1.5 x 10 <sup>9</sup> / L
Platelets	80 x 10 <sup>9</sup> / L

Reference: AML 19 trial protocol

Hypersensitivities/Allergies
<b>PRN antiemetics</b> ± Domperidone 10 mg PO QID ± Cyclizine 25 mg -50 mg IV / PO TDS

Agent	Round
Tretinoin	10 mg
Mitoxantrone	2 mg
<b>DOSE MODIFIED: No Yes</b>	

Day	Date	Time	Agent	Dose	Route	Instructions	Doctor	Nurse	Check	Start	Stop	
1			<b>Tretinoin (ATRA) 45 mg/m<sup>2</sup>/day</b> (split in 2 divided doses)	mg	BD	PO	Administered daily in 2 equally divided doses, from day 1 – 15. Chart on medication chart or outpatient prescription.					
			Ondansetron	8	mg	PO/IV	Give 1 hour prior to chemotherapy					
			<b>Mitoxantrone 10 mg/m<sup>2</sup></b>		mg	IV	In 50 mL sodium chloride 0.9 % over 30 minutes					
2			Ondansetron	8	mg	PO/IV	Give 1 hour prior to chemotherapy					
			<b>Mitoxantrone 10 mg/m<sup>2</sup></b>		mg	IV	In 50 mL sodium chloride 0.9 % over 30 minutes					

- You must read the protocol before prescribing this chemotherapy.

**Consultant:**  
**NZMC Reg No:**

**AML 19**  
**Acute Promyelocytic Leukaemia**  
**AIDA Consolidation cycle 2**

<b>Height</b>	cm
<b>Weight</b>	kg
<b>BSA</b>	m <sup>2</sup>

--

Day	Date	Time	Agent	Dose	Route	Instructions	Doctor	Nurse	Check	Start	Stop
3			Ondansetron	8 mg	PO/IV	Give 1 hour prior to chemotherapy					
			<b>Mitoxantrone 10 mg/m<sup>2</sup></b>	mg	IV	In 50 mL sodium chloride 0.9 % over 30 minutes					
4			Ondansetron	8 mg	PO/IV	Give 1 hour prior to chemotherapy					
			<b>Mitoxantrone 10 mg/m<sup>2</sup></b>	mg	IV	In 50 mL sodium chloride 0.9 % over 30 minutes					
5			Ondansetron	8 mg	PO/IV	Give 1 hour prior to chemotherapy					
			<b>Mitoxantrone 10 mg/m<sup>2</sup></b>	mg	IV	In 50 mL sodium chloride 0.9 % over 30 minutes					