

AML 19
Acute Promyelocytic Leukaemia
AIDA Consolidation cycle 3

Height	cm
Weight	kg
BSA	m ²

--

AML 19 patient trial number:

Destination:	BMTU
	HDW

	Limits
Date	
Neutrophils	1.5 x 10 ⁹ / L
Platelets	80 x 10 ⁹ / L

Reference: AML 19 trial protocol

Hypersensitivities/Allergies

PRN antiemetics
± Domperidone 10 mg PO QID
± Cyclizine 25 mg -50 mg IV / PO TDS

<u>Agent</u>	<u>Round</u>
Tretinoin	10 mg
Idarubicin	1 mg

DOSE MODIFIED: No Yes

Day	Date	Time	Agent	Dose	Route	Instructions	Doctor	Nurse	Check	Start	Stop		
1			Tretinoin (ATRA) 45 mg/m²/day (split in 2 divided doses)	mg	PO	Administered daily in 2 equally divided doses, from day 1 – 15. Chart on medication chart or outpatient prescription.							
			Ondansetron	8	mg							PO/IV	Give 1 hour prior to chemotherapy
			Idarubicin 12 mg/m²	mg	IV							In 50 mL sodium chloride 0.9 % over 20 minutes	

- You must read the protocol before prescribing this chemotherapy.

Consultant:
NZMC Reg No: