

Modified OPAL
without colaspase (l-asparaginase)

| | |
|---------------|----------------|
| Height | cm |
| Weight | kg |
| BSA | m ² |

| |
|--|
| |
|--|

| | | | | |
|--------------|------|-------|--------------|---------------|
| Destination: | BMTU | | Day 1 | Limits |
| | | Date: | | |
| | | ECHO | | |
| | | | | |

Hypersensitivities/Allergies

PRN antiemetics
± Domperidone 10 mg PO QID
± Cyclizine 25 mg - 50 mg IV / PO TDS

| | | |
|-----------------------|--------------|--------------|
| | <u>Agent</u> | <u>Round</u> |
| | Doxorubicin | 5 mg |
| | Vincristine | 0.2 mg |
| DOSE MODIFIED: | | |
| | No | Yes |

Reference: Lister, T. A, Whitehouse, J. M et al BMJ 1978:199 - 203

| Day | Date | Time | Agent | Dose | Route | Instructions | Doctor | Nurse | Check | Start | Stop |
|------|------|------|---|------|-------|--|--|-------|-------|-------|------|
| -1 | | | Methotrexate 12 mg | | IT | See separate intrathecal chemotherapy medication chart (C160016) | | | | | |
| 1-28 | | | Prednisone 40 mg/m² | | mg | PO | Prescribe on MedChart or outpatient prescription, then taper over 14 days. | | | | |
| 1 | | | Ondansetron | 8 | mg | PO/IV | Give 1 hour prior to chemotherapy | | | | |
| | | | Doxorubicin 30 mg/m² | | mg | IV | In 150 mL Sodium Chloride 0.9% over 15 min | | | | |
| | | | Vincristine 1.4 mg/m² (max 2mg) | | mg | IV | In 50mL sodium chloride 0.9% over 10 – 15 minutes | | | | |
| | | 2000 | Ondansetron | 8 | mg | PO/IV | | | | | |
| 2 | | | Ondansetron | 8 | mg | PO/IV | Give 1 hour prior to chemotherapy | | | | |
| | | | Doxorubicin 30 mg/m² | | mg | IV | In 150 mL Sodium Chloride 0.9% over 15 min | | | | |
| | | 2000 | Ondansetron | 8 | mg | PO/IV | | | | | |

Consultant:
NZMC Reg. No:

Modified OPAL
without colaspase (l-asparaginase)

| | |
|---------------|----------------|
| Height | cm |
| Weight | kg |
| BSA | m ² |

| |
|--|
| |
|--|

*Falling neutrophils < 0.5 x 10⁹/L and platelets < 50 x 10⁹/L require bone marrow and chemotherapy review before day 14 chemotherapy is given

| Day | Date | Time | Agent | Dose | Route | Instructions | Doctor | Nurse | Check | Start | Stop | |
|-----|------|------|--|------|-------|--------------|--|-------|-------|-------|------|--|
| 14 | | | Ondansetron | 8 | mg | PO/IV | Give 1 hour prior to chemotherapy | | | | | |
| | | | *Doxorubicin 30 mg/m² | | mg | IV | In 150 mL Sodium Chloride 0.9% over 15 min | | | | | |
| | | | *Vincristine 1.4 mg/m² (max 2mg) | | mg | IV | In 50mL sodium chloride 0.9% over 10 – 15 minutes | | | | | |
| | | 2000 | Ondansetron | 8 | mg | PO/IV | | | | | | |
| 21 | | | Ondansetron | 8 | mg | PO/IV | Give 1 hour prior to chemotherapy | | | | | |
| | | | Doxorubicin 30 mg/m² | | mg | IV | In 150 mL Sodium Chloride 0.9% over 15 min | | | | | |
| | | | Vincristine 1.4 mg/m² (max 2mg) | | mg | IV | In 50mL sodium chloride 0.9% over 10 – 15 minutes | | | | | |
| | | 2000 | Ondansetron | 8 | mg | PO/IV | | | | | | |
| 28 | | | Ondansetron | 8 | mg | PO/IV | Give 1 hour prior to chemotherapy | | | | | |
| | | | Doxorubicin 30 mg/m² | | mg | IV | In 150 mL Sodium Chloride 0.9% over 15 min | | | | | |
| | | | Vincristine 1.4 mg/m² (max 2mg) | | mg | IV | In 50mL sodium chloride 0.9% over 10 – 15 minutes | | | | | |
| | | 2000 | Ondansetron | 8 | mg | PO/IV | | | | | | |
| 29 | | | Prednisone | | mg | PO | Taper over 14 days | | | | | |
| 32 | | | Methotrexate 12 mg | | | IT | See separate intrathecal chemotherapy medication chart (C160016) | | | | | |