

GANCICLOVIR for Post ALLO SCT CMV prophylaxis						Deliver to: HDW					
Patient Category: Y J A P O 1 3 Z		OSP Group: 1 2 3 4 5 6 7		Doctors Name: NZMC Reg. No.:		Special Authority Number:					
						Antiemetics		Weight kg	Height cm		
								BSA m²			
								Dose Modified	No		
Give 3 days/week – Monday, Wednesday and Friday. Start at Day +21 if neutrophil count > 0.5 x10 ⁹ /L; or subsequently start once neutrophil count > 0.5 x10 ⁹ /L.						ADMINISTRATION USE					
DAY	DATE	TIME	MEDICATION	DOSE	ROUTE	INSTRUCTIONS	DR SIGN	GIVEN BY	CHECK	TIME	
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					

Chemotherapy rounding ganciclovir 10mg

- Note:**
- Stop at Day +84 if no GvHD; otherwise stop at Day +100
 - Stop if neutrophils drop to < 0.5 or platelets <20 x10⁹/L (Discuss with Consultant)
 - Monitor CMV viral load weekly (Monday), if this rises or CMV like illness develops therapeutic doses of ganciclovir may be needed. Discuss with consultant

Authorised by: Clinical Director Steve Gibbons Pharmacist Bevan Harden
 T:\Pictures\CDHB\Red_Book\Ganciclovir Post-SCT.doc

13 February 2008
 Ref: 2000

GANCICLOVIR for Post ALLO SCT CMV prophylaxis						Deliver to: HDW					
Patient Category: Y J A P O 1 3 Z		OSP Group: 1 2 3 4 5 6 7		Doctors Name: NZMC Reg. No.:		Special Authority Number:					
						Antiemetics		Weight kg	Height cm		
								BSA m²			
								Dose Modified	No		
Give 3 days/week – Monday, Wednesday and Friday. Do not give aciclovir concurrently. Start at Day +21 if neutrophil count > 0.5 x10 ⁹ /L; or subsequently start once neutrophil count > 0.5 x10 ⁹ /L.						ADMINISTRATION USE					
DAY	DATE	TIME	MEDICATION	DOSE	ROUTE	INSTRUCTIONS	DR SIGN	GIVEN BY	CHECK	TIME	
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					
			Ganciclovir 5mg/kg	mg	IV	In 100mL 0.9S over 1 hour					

Chemotherapy rounding ganciclovir 10mg

- Note:**
- Stop at Day +84 if no GvHD; otherwise stop at Day +100
 - Stop if neutrophils drop to < 0.5 or platelets <20 x10⁹/L (Discuss with Consultant)
 - Monitor CMV viral load weekly, if this rises or CMV like illness develops therapeutic doses of ganciclovir may be needed. Discuss with Consultant.

Authorised by: Clinical Director Steve Gibbons Pharmacist Bevan Harden
 T:\Pictures\CDHB\Red_Book\Ganciclovir Post-SCT.doc

13 February 2008
 Ref: 2000