

UKALL 14
Consolidation 1

Height	cm
Weight	kg
BSA	m ²

--

Cycle length:	21 days
Destination:	

CBC	Day 1	Limits
Date		
Hb		
Neuts		> 0.75 x 10 ⁹ /L
Plts		> 75 x 10 ⁹ /L

Hypersensitivities/Allergies

Antiemetics
Domperidone 10mg PO QID
± Cyclizine 50mg TDS PO/IV

Agent	Round
Cytarabine	10 mg
Etoposide	10 mg
Peg-Asparaginase	75 units (IU)

DOSE MODIFIED: Yes / No

Reference: UKALL 14 trial protocol –v5.0 – 20/07/12

Day	Date	Time	Agent	Dose	Route	Instructions	Doctor	Nurse	Check	Start	Stop
Continuous			Imatinib <i>(Philadelphia +ve patients only)</i>		mg PO	Continued for patients with Philadelphia positive disease . Continuous daily Imatinib, PO, aiming to escalate to 600mg, if tolerated. This should be continued until transplant wherever possible. Prescribe on regular medication chart or outpatient prescription.					
1			Methotrexate 12 mg		IT	See separate intrathecal chemotherapy medication chart (C160016)					
1			Ondansetron	8	mg PO	Give one hour prior to chemotherapy					
			Dexamethasone	8	mg PO	Give one hour prior to chemotherapy					
			Cytarabine 75 mg/m²		mg IV	Bolus					
			Etoposide 100 mg/m²		mg IV	In 500 mL sodium chloride 0.9 % over 1 hour					
			Ondansetron	8	mg PO						
2			Ondansetron	8	mg PO	Give one hour prior to chemotherapy					
			Dexamethasone	8	mg PO	Give one hour prior to chemotherapy					
			Cytarabine 75 mg/m²		mg IV	Bolus					
			Etoposide 100 mg/m²		mg IV	In 500 mL sodium chloride 0.9 % over 1 hour					
			Ondansetron	8	mg PO						

▪ Timing of intrathecal therapy can be moved +/- 3 days to allow administration on specific lists as per local and national guidelines

Consultant:
NZMC Reg. No:

**UKALL 14
Consolidation 1**

Height	cm
Weight	kg
BSA	m ²

--

Day	Date	Time	Agent	Dose	Route	Instructions	Doctor	Nurse	Check	Start	Stop	
3			Ondansetron	8 mg	PO	Give one hour prior to chemotherapy						
			Dexamethasone	8 mg	PO	Give one hour prior to chemotherapy						
			Cytarabine 75 mg/m²		mg	IV	Bolus					
			Etoposide 100 mg/m²		mg	IV	In 500 mL sodium chloride 0.9 % over 1 hour					
			Ondansetron	8 mg	PO							
4			Ondansetron	8 mg	PO	Give one hour prior to chemotherapy						
			Dexamethasone	8 mg	PO	Give one hour prior to chemotherapy						
			Cytarabine 75 mg/m²		mg	IV	Bolus					
			Etoposide 100 mg/m²		mg	IV	In 500 mL sodium chloride 0.9 % over 1 hour					
			Ondansetron	8 mg	PO							
5			Ondansetron	8 mg	PO	Give one hour prior to chemotherapy						
			Dexamethasone	8 mg	PO	Give one hour prior to chemotherapy						
			Peg-Asparaginase 1000 units/m²		units	IV	In 100 mL sodium chloride 0.9 % over at least 1 hour					
			Cytarabine 75 mg/m²		mg	IV	Bolus					
			Etoposide 100 mg/m²		mg	IV	In 500 mL sodium chloride 0.9 % over 1 hour					
			Ondansetron	8 mg	PO							
6			Dexamethasone	8 mg	PO							
7			Dexamethasone	8 mg	PO							

- Check Antithrombin III and Fibrinogen every 2 days while receiving Peg-Asparaginase
- Consolidation 2 to commence 3 weeks from the first day of consolidation 1 or when neutrophils > 0.75 X 10⁹/L and platelets >75 X 10⁹/L