

Non Hodgkin Lymphoma (NHL)
Relapsed Lymphoma
Vinorelbine & gemcitabine

| | |
|---------------|----------------|
| Height | cm |
| Weight | kg |
| BSA | m ² |

| |
|--|
| |
|--|

| | |
|---------------|---------|
| Cycle length: | 21 days |
| Cycle number: | |
| Destination: | |

| CBC | Day 1 | Limits |
|-------|-------|----------------------------|
| Date | | |
| Hb | | |
| Neuts | | > 0.5 x 10 ⁹ /L |
| Plts | | > 50 x 10 ⁹ /L |

Reference: Pasricha et al, Cancer 2008; 113:3192-8

Hypersensitivities/Allergies

PRN antiemetics
Domperidone 10 mg PO QID PRN
± Cyclizine 50 mg TDS PO/IV PRN

| Agent | Round |
|-------------|-------|
| Vinorelbine | 1 mg |
| Gemcitabine | 10 mg |

DOSE MODIFIED: No Yes

| Day | Date | Time | Agent | Dose | Route | Instructions | Doctor | Nurse | Check | Start | Stop | |
|-----|------|------|--|------------|-------|------------------------------|---|-------|-------|-------|------|--|
| 1 | | | Ondansetron | 8 mg | PO | 1 hour prior to chemotherapy | | | | | | |
| | | | Sodium chloride 0.9% | 100-500 mL | IV | For flushing | | | | | | |
| | | | Vinorelbine 25 mg/m² | | mg | IV | In 50 ml sodium chloride 0.9%, free run over 10 minutes | | | | | |
| | | | Gemcitabine 1000 mg/m² | | mg | IV | In 150 mL sodium chloride 0.9% over 30 minutes | | | | | |

Consultant:
NZMC Reg. No:

Special Authority:
Pegfilgrastim:

Consider the following supportive medications – see outpatient prescription template in Red Book:

- **Allopurinol 100mg OR 300mg PO once daily for 4 weeks**
- **Valaciclovir 500mg PO once OR twice daily**
- **Sulfamethoxazole + trimethoprim 480mg PO once daily**

Non Hodgkin Lymphoma (NHL)
Relapsed Lymphoma
Vinorelbine & gemcitabine

| | |
|---------------|----------------|
| Height | cm |
| Weight | kg |
| BSA | m ² |

| |
|--|
| |
|--|

| Day | Date | Time | Agent | Dose | Route | Instructions | Doctor | Nurse | Check | Start | Stop |
|-----|------|------|--|----------|-----------|---|--------|-------|-------|-------|------|
| 8 | | | Ondansetron | 8 | mg PO | 1 hour prior to chemotherapy | | | | | |
| | | | Sodium chloride 0.9% | 100-500 | mL IV | For flushing | | | | | |
| | | | Vinorelbine 25 mg/m² | | mg IV | In 50 ml sodium chloride 0.9%, free run over 10 minutes | | | | | |
| | | | Gemcitabine 1000 mg/m² | | mg IV | In 150 mL sodium chloride 0.9% over 30 minutes | | | | | |
| 9 | | | Pegfilgrastim | 6 | mg subcut | Inject subcutaneously at least 24 hours following last dose of chemotherapy. <i>Prescribe on an outpatient prescription. Special authority required.</i> | | | | | |